



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at ScienceDirect

Atherosclerosis

journal homepage: www.elsevier.com/locate/atherosclerosis

Associations of birth weight with serum long chain polyunsaturated fatty acids in adolescents; the HELENA study

Idoia Labayen^{a,b,*}, Luis A. Moreno^{b,1}, Jonatan R. Ruiz^{c,d,1}, Francisco B. Ortega^{d,e,1}, Michael Sjoström^{d,1}, Inge Huybrechts^{f,1}, Marcela Gonzalez-Gross^{g,1}, André Spinneker^{h,1}, Stefaan De Henauw^{f,1}, Yannis Manios^{i,1}, Dénés Molnár^{j,1}, Catherine Leclercq^{k,1}, Frédéric Gottrand^{l,1}, on behalf of the HELENA study group¹

^a Department of Nutrition and Food Science, University of the Basque Country, Vitoria, Spain

^b GENUD (Growth, Exercise, Nutrition and Development) Research Group, Escuela Universitaria de Ciencias de la Salud, Universidad de Zaragoza, 50009 Zaragoza, Spain

^c Department of Physical Education and Sport, School of Physical Activity and Sport Sciences, University of Granada, Granada, Spain

^d Unit for Preventive Nutrition, Department of Biosciences and Nutrition at NOVUM, Karolinska Institutet, SE-17177 Huddinge, Sweden

^e Department of Medical Physiology, School of Medicine, University of Granada, 18014 Granada, Spain

^f Department of Public Health, Faculty of Medicine and Health Sciences, Ghent University, Ghent, Belgium

^g Faculty of Physical Activity and Sport Sciences, University Polytechnic of Madrid, Madrid, Spain

^h Department of Nutrition and Food Science, University of Bonn, Germany

ⁱ Department of Nutrition and Dietetics, Harakopio University, Athens, Greece

^j Department of Paediatrics, Medical Faculty, Pecs, Hungary

^k INRAN (National Research Institute on Food and Nutrition), Via Ardeatina, 546100176 Roma, Italy

^l INSERM U995, IFR114, Faculté de Médecine, Université Droit et Santé de Lille 2, Lille, France

ARTICLE INFO

Article history:

Received 8 February 2011

Received in revised form 14 March 2011

Accepted 25 March 2011

Available online 5 April 2011

Keywords:

Birth weight

Phospholipids

Long chain polyunsaturated fatty acids

Adolescents

HELENA study

ABSTRACT

Objectives: Nutritional factors in early life may have long-term physiologic effects in humans. Eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) play important roles in protecting against cardiovascular disease (CVD) risk. Our aim was to examine the association of birth weight (BW) with serum long chain polyunsaturated fatty acids (LCPUFA) profile in adolescents.

Subjects and methods: A total of 772 European adolescents (56.3% females) aged 14.7 ± 1.2 years were included in this study. Information on BW and gestational age was obtained from parental records. DHA, EPA and arachidonic acid (AA) concentrations were measured in serum phospholipids. Alfa-linolenic (ALA), linoleic (LA), AA, EPA and DHA intakes assessed by a computer based 24 h dietary recall. Gender, gestational age, pubertal status, body mass index, center and total energy and LCPUFA intakes were used as confounders in all the analyses.

Results: BW was significantly associated with serum DHA and EPA (both adjusted $P < 0.05$) independently of potential confounders including their main dietetic source. We did not observe any significant relationship between BW and serum AA levels.

Conclusions: Our findings suggest that early metabolic changes, as a result from prenatal environmental influences, could affect long chain polyunsaturated fatty acid metabolism later in life. These results may contribute to explain the relationship between early nutrition and growth and later metabolic disorders as CVD.

© 2011 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

There is now compelling evidence that low birth weight (BW) is associated with atherosclerosis and cardiovascular disease (CVD)

* Corresponding author at: Nutrición y Bromatología, Facultad de Farmacia, Universidad del País Vasco, Paseo de la Universidad, 7, 01006 Vitoria-Gasteiz, Spain. Tel.: +34 945014346; fax: +34 945130756.

E-mail address: idoia.labayen@ehu.es (I. Labayen).

¹ On behalf of the HELENA study group please see [Appendix A](#).

risk factors such as obesity, hypertension and type 2 diabetes mellitus later in life [1–6]. Environmental and genetic factors, as well as their interactions, seem to partially explain these associations [7,8].

The role of n-3 long chain polyunsaturated fatty acids (LCPUFA) in human health and disease has received special attention during recent decades. Eicosapentaenoic acid (C20:5n-3, EPA) and docosahexaenoic acid (C22:6n-3, DHA) play important roles in human health through various mechanisms, including effects against inflammation, platelet aggregation, hypertension and hyperlipidemia [9]. Thus, an inverse association between EPA and DHA

content of blood phospholipids and CVD risk and subclinical atherosclerosis is a consistent finding in observational studies [10–12]. Furthermore, serum DHA levels have been inversely associated with common carotid artery intima media thickness in subjects with primary dyslipemia [13]. Arachidonic acid (C20:4n-6, AA) is the precursor of important molecules involved in inflammation (e.g. eicosanoids), and is thought to play a role in the atherosclerotic process [14,15]. High concentrations of AA in adipose tissue have been associated with a greater risk of myocardial infarction, which suggests a pro-atherosclerotic role of excess of AA [16,17].

Tissue LCPUFA levels are mainly determined by dietary intake and endogenous synthesis, via the successive elongation and desaturation of dietary polyunsaturated fatty acids precursors, linoleic-acid (C18:2n-6, LA) and α -linolenic acid (C18:3n-3, ALA). ALA is converted to EPA and then to DHA, whereas LA is converted to AA through enzymatic chain elongation and desaturation. Although LCPUFA intakes and metabolic turnover are likely to be the main predictors of DHA and EPA status [18], there are also other factors that influence DHA and EPA levels including age, nutritional status or genetic background [19–22]. Serum phospholipid DHA and EPA are effective biomarkers of DHA and EPA status, respectively [23].

Longitudinal studies suggested that maternal nutritional status during ovum maturation, or even before conception, has a greater effect on the neonatal fatty acid profile than maternal nutrition during the last 2 trimesters of pregnancy [24,25]. It has also been shown that neonatal LCPUFA concentrations at birth seem to significantly affect the postnatal LCPUFA content in blood phospholipids [26]. Several studies suggested that undernutrition and suboptimal growth in pregnancy could change the metabolism of cholesterol and triglycerides [8,27,28]. Information is lacking on the effect of BW on serum LCPUFA concentrations later in life.

The present study aimed to examine the association of BW with serum AA, EPA and DHA levels in European adolescents participating in the Healthy Lifestyle in Europe by Nutrition in Adolescence Study (HELENA).

2. Subjects and methods

2.1. Subjects

The recruitment and phenotyping of the adolescents participating in the HELENA cross-sectional study (“Healthy Lifestyle in Europe by Nutrition in Adolescence”, www.helenastudy.com) have been described previously (18). Briefly, a total of 3865 adolescents were recruited between 2006 and 2007. Data were collected in a total of 10 centers from 9 European countries. Subjects were randomly selected from schools by using a proportional cluster sampling methodology and taking age into account. One third of the classes were randomly selected for blood collection; this resulted in a total of 1155 blood samples.

Data were collected on a detailed case report form, in accordance with standardized procedures. In each center, trained researchers carried out complete physical examinations, including weight, height and blood pressure measurements. The protocol was approved by the appropriate investigational review board for each investigating center. Written, informed consent was obtained from each adolescent and both of his/her parents or legal representatives. Participation in the study was voluntary.

For the purpose of this study, adolescents with at least data on serum AA, EPA and DHA levels, body mass index (BMI) and neonatal data were included in the analysis. To be born at term (95.5%) was an additional inclusion criterion. Based on these criteria, the final study sample comprised 772 adolescents (56.3% females). The final

sample did not differ in key characteristics (i.e., BW and age) from the original sample (all $P > 0.1$).

2.2. Neonatal data

A questionnaire was developed for parents to collect information on the adolescents' BW and gestational age [29]. Parents were specifically asked to recall this information from the health booklets of their son/daughter. This questionnaire was sent to the parents together with the study information letter and consent form, and collected at school on the first day of the examinations. If information from the parental questionnaire was lacking, the local investigators were advised to send the questionnaire to the parents again to obtain the required information.

2.3. Physical examination

Harmonization and standardization of anthropometric measurements used to assess body composition in the HELENA Study were strictly controlled and have been previously described [30]. Weight (Type SECA 861; range, 0.05–130 kg; precision, 0.05 kg), and height (Type SECA 225; range, 60–200 cm; precision, 1 mm) were measured. Thereafter, BMI was calculated. Identification of pubertal status (stages I–V) was assessed by a medical doctor according to Tanner and Whitehouse [31].

2.4. Serum phospholipid LCPUFA analysis

Blood samples were drawn at school according to standardized collection protocol (after 10-h overnight fast)[32]. Samples were centrifuged directly at school (3500 rpm for 15 min at room temperature) and serum stored at -20°C . Serum samples (1 mL) were defrosted at room temperature and 50 μL internal standard (1,2-dipentadecanoyl-sn-glycero-3-phosphocholine, Sigma–Aldrich, Deisenhofen, Germany) were added. A methanol/chloroform solution was used to extract proteins and transfer lipid classes into a non-aqueous solution according to the modified methods of Folch [33]. Briefly, 1 mL of methanol and 2 mL of chloroform were added to the sample. The solution was centrifuged (2000 rpm, 10 min, 4°C). The upper layer was removed via aspiration and the interphase by filtration. The filtrate was evaporated to dryness under a N_2 flow and the residue was dissolved in chloroform. After each step, N_2 was blown over the samples to avoid fatty acid (FA) oxidation. The serum phospholipid fraction was separated by thin-layer chromatography according to the method of Christophe and Matthijs [34]. The phospholipid band was scraped off and the FA were converted into their methyl esters by transesterification using 2 mL of a mixture of methanol/hydrochloric acid as methylating solution for 4 h at 95°C . After cooling (-20°C , 10 min), distilled water was added. The methyl esters were extracted with petroleum ether (bp 40 – 60°C) and evaporated to dryness under a N_2 flow. The residue was redissolved in heptane and analyzed. The FA methyl esters of the phospholipid fraction were analyzed using a temperature-programmed capillary gas chromatograph (Model 3900, Varian GmbH, Darmstadt, Germany) with a $30\text{ m} \times 0.25\text{ mm} \times 0.25\text{ }\mu\text{m}$ polyethylene glycol column (Zebron ZB-WAXplus, Phenomenex Ltd., Aschaffenburg, Germany). Helium was used as carrier gas with a constant flow of 4 mL/min, the split ratio was set to 1:20. Injector and flame ionization detector temperatures were 200°C and 250°C , respectively. The oven temperature was programmed as follows: the initial temperature (100°C) was held for 2 min and increased to 210°C at rates of $25^{\circ}\text{C}/\text{min}$, followed by $0.5^{\circ}\text{C}/\text{min}$, $1.0^{\circ}\text{C}/\text{min}$, and $5.0^{\circ}\text{C}/\text{min}$, respectively. After reaching 210°C , the temperature was increased to 240°C at a rate of $10^{\circ}\text{C}/\text{min}$. The final temperature of 240°C was held for 3 min. Peaks of interest were

Table 1
Characteristics of the study sample.

	Males (n = 337)	Females (n = 435)	All (n = 772)
Age (year)	14.7 ± 1.2	14.7 ± 1.1	14.7 ± 1.2
Birth weight (kg)	3.4 ± 0.6	3.3 ± 0.6	3.3 ± 0.6
Body mass index (kg/m ²)	21.2 ± 3.9	21.3 ± 3.5	21.2 ± 3.7
AA (μmol/L)	343.4 ± 89.4	363.4 ± 98.7	354.6 ± 95.1
EPA (μmol/L)	18.6 ± 13.6	19.1 ± 14.1	18.9 ± 13.9
DHA (μmol/L)	94.3 ± 40.0	112.0 ± 40.7	104.2 ± 41.5
Total energy intake (kcal/day)	2570 ± 895	1892 ± 520	2189 ± 784
LA intake (mg/day)	10800 ± 5641	8260 ± 3221	9373 ± 4618
ALA intake (mg/day)	1697 ± 1162	1255 ± 590	1449 ± 913
AA intake (mg/day) [*]	321 ± 182	251 ± 152	282 ± 170
EPA intake (mg/day) [*]	70 ± 61	63 ± 54	66 ± 58
DHA intake (mg/day) [*]	166 ± 142	159 ± 140	162 ± 141

Data are expressed as means ± standard deviation unless otherwise stated. AA: arachidonic acid; ALA: alfa-linolenic fatty acid; DHA: docosahexaenoic fatty acid; EPA: eicosapentaenoic fatty acid; LA: linoleic fatty acid.

^{*} N = 511 adolescents (224 males and 287 females).

identified by comparison with authentic FA methyl ester standards (Sigma–Aldrich, Deisenhofen, Germany). The absolute FA composition was expressed in μmol/L. The coefficients of variations for the analyses were below 4.4% for all FA.

For the present study, the following fatty acids were relevant: AA; EPA; and DHA. Other measured fatty acids were not considered because they were beyond the scope of our study.

2.5. Dietary intake assessment

To obtain dietary intake data, the HELENA-DIAT 24-h dietary recall software was used. This 24-h recall assessment tool is based on six meal occasions referring to the day before the interview. The adolescents completed the questionnaire during school time, after dieticians/researchers instructed them on how to fill in this 24-h recall as accurately as possible. The participants were allowed to ask questions and assistance and after completion, the recall was checked for completeness. Every participant was asked to fill in the HELENA-DIAT on arbitrary days, twice in a time-span of 2 weeks. Since the questionnaire was filled in during school time, no data could be collected about the dietary intake on Fridays and Saturdays. A validation study by Vereecken et al. [35] indicated that the YANA-C, a former version of the HELENA-DIAT, showed good agreement with an interviewer-administered YANA-C interview. The HELENA-DIAT tool has been indicated as a good method to collect detailed dietary information from adolescents and was received well by the study participants [35]. Furthermore, a repeated 24-h recall was selected as the most suitable method to get population means and distributions by the European Consumption Survey Method (EFCOSUM) project [36]. To calculate energy and nutrient intake, data of the HELENA-DIAT was linked to the German Food Code and Nutrient Data Base (BLS (Bundeslebensmittelschlüssel), version II.3.1, 2005).

Table 2
Multiple regression standardized coefficients (β) and standard errors (SE) examining the association of birth weight with serum long chain polyunsaturated fatty acid concentrations.

	Unadjusted (n = 772)			Model 2 (n = 772)			Model 3 (n = 511)		
	β	SE	P	β	SE	P	β	SE	P
AA (μmol/L) ^a	−0.003	0.016	0.874	0.021	0.016	0.199	0.028	0.019	0.140 ^b
DHA (μmol/L) ^a	0.051	0.024	0.035	0.057	0.023	0.013	0.054	0.026	0.039 ^c
EPA (μmol/L) ^a	0.156	0.041	<0.001	0.076	0.037	0.039	0.088	0.041	0.031 ^c

AA: arachidonic acid; DHA: docosahexaenoic acid; EPA: eicosapentaenoic acid. Model 2 controls for pubertal status, sex, study location and BMI. Model 3: includes model 1 and additionally controls for total energy intake and the corresponding polyunsaturated fatty acid dietetic source.

^a Analysis was performed with log-transformed data.

^b Adjusted for linoleic and arachidonic acids intake.

^c Adjusted for alfa-linolenic, docosahexaenoic and eicosapentaenoic acids intake.

In addition, the adolescents completed a brief Food Frequency Questionnaire (FFQ) asking for the frequency in consumption of different food items (e.g. fish consumption). The frequency categories included the options 'never', 'less than once per week', 'once per week', '2 to 4 times per week', '5 to 6 times per week', 'once per day' and 'more times per day'. Usual food intakes were estimated for each adolescent based on the 24-h recalls and the FFQ-information by using the Multiple Source Method (MSM) (<https://nugo.dife.de/msm/>). The MSM calculates dietary intake for individuals first and then constructs the population distribution based on the individual data. With this method dietary data was corrected for between and within person variability.

For the purpose of this study we used as confounders ALA, DHA and EPA as the main dietetic sources of plasma DHA and EPA levels, and LA and AA as the main dietetic sources of plasma AA levels, respectively. We also recorded total energy intake (kcal/day).

3. Statistical analysis

Statistical analyses were performed using "Statistical Package for the Social Sciences (SPSS)" software 17.0 (SPSS, Chicago, IL), and the threshold for statistical significance was set at $P < 0.05$. Physical and biochemical characteristics of the study sample by sex are presented as means and standard deviation, unless otherwise stated. Variables with skewed distribution, i.e., AA, EPA, and DHA levels, were log transformed to obtain a more symmetrical distribution.

Regression analysis was used to examine the association between BW and serum phospholipid LCPUFA levels. These relationships were analyzed in three separate regression models. The unadjusted model included the predictor (BW) and the dependent variables (serum AA, EPA and DHA levels). Model 2 included BW and the dependent variable adjusted for gestational age, pubertal

status, sex, center (entered as dummy variable) and BMI. For model 3, analyses were additionally controlled for dietary intake variables. Likewise, regression analysis examining the association between BW and serum AA levels was controlled for LA, AA and energy intake variables, and the relationships between BW and serum EPA and DHA levels were additionally adjusted for ALA, EPA, DHA, and energy intake variables:

4. Results

The descriptive characteristics of the participants are presented in Table 1.

Table 2 presents linear regression statistics showing the observed change in mean serum phospholipids LCPUFA levels per kilogram increase in BW in adolescents. BW was positively associated with serum EPA ($P < 0.05$) and DHA ($P < 0.001$) levels in the unadjusted model and these relationships remain statistically significant after additional adjustment for sex, gestational age, BMI, pubertal status and center (model 2, Table 2). Further controlling for total energy, ALA, EPA and DHA intakes did not substantially change the outcome (model 3, Table 2).

BW was not significantly associated with serum AA levels in any of the studied regression models (Table 2).

5. Discussion

The findings of the present study show significant associations between BW and serum DHA and EPA levels in adolescents independently of potential confounders, including ALA, DHA and EPA intakes. To our knowledge, there are no previous studies examining the association of BW on later serum LCPUFA profile, which hamper between studies comparisons.

It is biologically plausible that environmental intrauterine factors, such as insufficient fetus nutrition and growth, may partially program the activity and/or expression of the enzymes required for the metabolic endogenous synthesis of these LCPUFA. In utero challenges, such as relative under-provision of nutrients or placental insufficiency, can result in the long-term programming of individual metabolism, tissues and whole organ systems, with adverse consequences for their function in later life. Alternatively, the familial resemblance of dietary habits between the mothers and their children could be responsible for long term LCPUFA tracking. Likewise, to obtain LCPUFA the fetus depends on placental transfer and, thus, on the LCPUFA status of the mother. Indeed, an adverse maternal fatty acid profile early in pregnancy has been associated with reduced fetal growth [35]. Previous reports [35–38] estimated that low maternal serum concentrations of n-3 PUFA increased the risk of being small for gestational age in 40–50%, and it was associated with decreased BW on an average of 50–60 g.

It could also be argued that the main determinants of serum phospholipid fatty acid profile in the short term are dietary habits, yet the results were consistent even after adjustment for DHA and EPA intake, as well as for their precursor ALA intake. It has been suggested that there are other factors influencing serum DHA and EPA levels. Previous studies reported considerable variability among individuals in the conversion rates of ALA and LA on their respective n-3 and n-6 metabolites, even when the subjects followed similar background diets [22,39]. Other reports showed that either obesity or under-nutrition in children can alter fatty acid composition in serum and erythrocyte membrane lipids [21,40,41]. Nevertheless, our results were consistent after controlling for dietary sources, yet caution should be paid because of the instrument used to assess dietary intake. Moreover, we do not know whether serum LCPUFA levels will maintain stable over and after the sexual development

and how changes in physical activity and dietary patterns could affect this apparent programming effect of BW.

In conclusion, type of dietary fatty acid intake plays an important role in modulating LCPUFA metabolism. It is also possible, however, that early metabolic changes, such as prenatal environmental influences, affect LCPUFA metabolism later in life. The relationship between BW and serum EPA and DHA levels may contribute to explain the relationship between low BW and later metabolic disorders such as atherosclerosis or CVD.

Acknowledgments

We acknowledge all participating children and adolescents, as well as their parents and teachers for their collaboration.

Financial support: The HELENA study was carried out with the financial support of the European Community Sixth RTD Framework Programme (Contract FOODCT-2005-007034). This manuscript was also partially supported by Cognis GmbH, by the Swedish Council for Working Life and Social Research (FAS), and by grants from the Spanish Ministry of Education (EX-EX-2008-0641, the Spanish Ministry of Health: Maternal, Child Health and Development Network (number RD08/0072), the Spanish Ministry of Science and Innovation (RYC-2010-05957), and the University of the Basque Country. The writing group takes sole responsibility for the content of this article. The content of this paper reflects only the authors' views, and the European Community is not liable for any use that may be made of the information contained therein. None of the authors had any personal or financial conflict of interest.

Appendix A. HELENA Study Group

Co-ordinator: Luis A. Moreno.

Core Group members: Luis A. Moreno, Frédéric Gottrand, Stefaan De Henauw, Marcela González-Gross, Chantal Gilbert.

Steering Committee: Anthony Kafatos (President), Luis A. Moreno, Christian Libersa, Stefaan De Henauw, Sara Castelló, Frédéric Gottrand, Mathilde Kersting, Michael Sjöstrom, Dénes Molnár, Marcela González-Gross, Jean Dallongeville, Chantal Gilbert, Gunnar Hall, Lea Maes, Luca Scalfi.

Project Manager: Pilar Meléndez.

1. Universidad de Zaragoza (Spain)

Luis A. Moreno, Jesús Fleta, José A. Casajús, Gerardo Rodríguez, Concepción Tomás, María I. Mesana, Germán Vicente-Rodríguez, Adoración Villarroya, Carlos M. Gil, Ignacio Ara, Juan Revenga, Carmen Lachen, Juan Fernández Alvira, Gloria Bueno, Aurora Lázaro, Olga Bueno, Juan F. León, Jesús M^a Garagorri, Manuel Bueno, Juan Pablo Rey López, Iris Iglesia, Paula Velasco, Silvia Bel, Theodora Mouratidou.

2. Consejo Superior de Investigaciones Científicas (Spain)

Ascensión Marcos, Julia Wärnberg, Esther Nova, Sonia Gómez, Ligia Esperanza Díaz, Javier Romeo, Ana Veses, Belén Zapatera, Tamara Pozo, David Martínez.

3. Université de Lille 2 (France)

Laurent Beghin, Christian Libersa, Frédéric Gottrand, Catalina Iliescu, Juliana Von Berlepsch.

4. Research Institute of Child Nutrition Dortmund, Rheinische Friedrich-Wilhelms-Universität Bonn (Germany)

Mathilde Kersting, Wolfgang Sichert-Hellert, Ellen Koeppen.

5. Pécsi Tudományegyetem (University of Pécs) (Hungary)

Dénes Molnár, Eva Erhardt, Katalin Csernus, Katalin Török, Szilvia Bokor, Mrs. Angster, Enikő Nagy, Orsolya Kovács, Judit Répasi.

6. University of Crete School of Medicine (Greece)

Anthony Kafatos, Caroline Codrington, María Plada, Angeliki Papadaki, Katerina Sarri, Anna Viskadourou, Christos Hatzis, Michael Kiriakakis, George Tsibinos, Constantine Vardavas, Manolis Sbokos, Eva Protyeraki, Maria Fasoulaki.

7. Institut für Ernährungs- und Lebensmittelwissenschaften – Ernährungphysiologie. Rheinische Friedrich Wilhelms Universität (Germany)

Peter Stehle, Klaus Pietrzik, Marcela González-Gross, Christina Breidenassel, Andre Spinneker, Jasmin Al-Tahan, Miriam Segoviano, Anke Berchtold, Christine Bierschbach, Erika Blatzheim, Adelheid Schuch, Petra Pickert.

8. University of Granada (Spain)

Manuel J. Castillo, Ángel Gutiérrez, Francisco B Ortega, Jonatan R Ruiz, Enrique G Artero, Vanesa España, David Jiménez-Pavón, Palma Chillón, Cristóbal Sánchez-Muñoz, Magdalena Cuenca

9. Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione (Italy)

Davide Arcella, Elena Azzini, Emma Barrison, Noemi Bevilacqua, Pasquale Buonocore, Giovina Catasta, Laura Censi, Donatella Ciarapica, Paola D'Acapito, Marika Ferrari, Myriam Galfo, Cinzia Le Donne, Catherine Leclercq, Giuseppe Maiani, Beatrice Mauro, Lorenza Mistura, Antonella Pasquali, Raffaella Piccinelli, Angela Polito, Raffaella Spada, Stefania Sette, Maria Zaccaria.

10. University of Napoli “Federico II” Dept of Food Science (Italy)

Luca Scalfi, Paola Vitaglione, Concetta Montagnese.

11. Ghent University (Belgium)

Ilse De Bourdeaudhuij, Stefaan De Henauw, Tineke De Vriendt, Lea Maes, Christophe Matthys, Carine Vereecken, Mieke de Maeyer, Charlene Ottevaere, Inge Huybrechts.

12. Medical University of Vienna (Austria)

Kurt Widhalm, Katharina Philipp, Sabine Dietrich, Birgit Kubelka Marion Boriss-Riedl.

13. Harokopio University (Greece)

Yannis Manios, Eva Grammatikaki, Zoi Bouloubasi, Tina Louisa Cook, Sofia Eleutheriou, Orsalia Consta, George Moschos, Ioanna Katsaroli, George Kraniou, Stalo Papoutsou, Despoina Keke, Ioanna Petraki, Elena Bellou, Sofia Tanagra, Kostalena Kallianoti, Dionysia Argyropoulou, Katerina Konadaki, Stamatoula Tsikrika, Christos Karaiskos.

14. Institut Pasteur de Lille (France)

Jean Dallongeville, Aline Meirhaeghe.

15. Karolinska Institutet (Sweden)

Michael Sjöström, Jonatan R Ruiz, Francisco B. Ortega, María Hagströmer, Anita Hurtig Wennlöf, Lena Hallström, Emma Patterson, Lydia Kwak, Julia Wärnberg, Nico Rizzo. **Asociación de Investigación de la Industria Agroalimentaria (Spain)**

Jackie Sánchez-Molero, Sara Castelló, Elena Picó, Maite Navarro, Blanca Viadel, José Enrique Carreres, Gema Merino, Rosa Sanjuán, María Lorente, María José Sánchez.

16. Campden BRI (United Kingdom)

Chantal Gilbert, Sarah Thomas, Elaine Allchurch, Peter Burgess.

17. SIK – Institutet foer Livsmedel och Bioteknik (Sweden)

Gunnar Hall, Annika Astrom, Anna Sverkéén, Agneta Broberg.

18. Meurice Recherche & Development asbl (Belgium)

Annick Masson, Claire Lehoux, Pascal Brabant, Philippe Pate, Laurence Fontaine.

19. Campden & Chorleywood Food Development Institute (Hungary)

Andras Sebok, Tunde Kuti, Adrienn Hegyi.

20. Productos Aditivos SA (Spain)

Cristina Maldonado, Ana Lorente.

21. Cárnicas Serrano SL (Spain)

Emilio García.

22. Cederroth International AB (Sweden)

Holger von Fircks, Marianne Lilja Hallberg, Maria Messerer

23. Lantmännen Food R&D (Sweden)

Mats Larsson, Helena Fredriksson, Viola Adamsson, Ingmar Börjesson.

24. European Food Information Council (Belgium)

Laura Fernández, Laura Smillie, Josephine Wills.

25. Universidad Politécnica de Madrid (Spain)

Marcela González-Gross, Agustín Meléndez, Pedro J. Benito, Javier Calderón, David Jiménez-Pavón, Jara Valtueña, Paloma Navarro, Alejandro Urzanqui, Ulrike Albers, Raquel Pedrero, Juan José Gómez Lorente.

References

- [1] Labayen I, Ortega FB, Sjöström M, Ruiz JR. Early life origins of low-grade inflammation and atherosclerosis risk in children and adolescents. *J Pediatr* 2009;155:673–7.
- [2] Ramadhani MK, Grobbee DE, Bots ML, Castro Cabezas M, Vos LE, Oren A, et al. Lower birth weight predicts metabolic syndrome in young adults: the Atherosclerosis Risk in Young Adults (ARYA)-study. *Atherosclerosis* 2006;184:21–7.
- [3] Labayen I, Moreno LA, Blay MG, Blay VA, Mesana MI, Gonzalez-Gross M, et al. Early programming of body composition and fat distribution in adolescents. *J Nutr* 2006;136:147–52.
- [4] Labayen I, Moreno LA, Ruiz JR, Gonzalez-Gross M, Warnberg J, Breidenassel C, et al. Small Birth Weight and Later Body Composition and Fat Distribution in Adolescents. The AVENA study. *Obesity* 2008;16:1680–6.
- [5] Labayen I, Ruiz JR, Vicente-Rodríguez G, Turck D, Rodríguez G, Meirhaeghe A, et al. Early life programming of abdominal adiposity in adolescents. The HELENA study. *Diabetes Care* 2009;32:2120–2.
- [6] Ortega FB, Labayen I, Ruiz JR, Martín-Matillas M, Vicente-Rodríguez G, Redondo C, et al. The AVENA study: are muscular and cardiovascular fitness partially programmed at birth? role of body composition. *J Pediatr* 2008;154:61–6.
- [7] Labayen I, Moreno LA, Martí A, Gonzalez-Lamuno D, Warnberg J, Ortega FB, et al. Effect of the Ala12 allele in the PPARgamma-2 gene on the relationship between birth weight and body composition in adolescents: the AVENA study. *Pediatr Res* 2007;62:615–9.
- [8] Ruiz JR, Labayen I, Ortega FB, Moreno LA, Gonzalez-Lamuno D, Martí A, et al. Birth weight and blood lipid levels in Spanish adolescents: influence of selected APOE APOC3 and PPARgamma2 gene polymorphisms. The AVENA study. *BMC Med Genet* 2008;9:98.
- [9] Riediger ND, Othman RA, Suh M, Moghadasian MH. A systemic review of the roles of n-3 fatty acids in health and disease. *J Am Diet Assoc* 2009;109:668–79.
- [10] Lindqvist HM, Sandberg AS, Fagerberg B, Hulthe J. Plasma phospholipid EPA and DHA in relation to atherosclerosis in 61-year-old men. *Atherosclerosis* 2009;205:574–8.
- [11] Albert CM, Campos H, Stampfer MJ, Ridker PM, Manson JE, Willett WC, et al. Blood levels of long-chain n-3 fatty acids and the risk of sudden death. *N Engl J Med* 2002;346:1113–8.
- [12] Simon JA, Hodgkins ML, Browner WS, Neuhaus JM, Bernert Jr JT, Hulley SB. Serum fatty acids and the risk of coronary heart disease. *Am J Epidemiol* 1995;142:469–76.
- [13] Sala-Vila A, Cofan M, Perez-Heras A, Nunez I, Gilabert R, Junyent M, et al. Fatty acids in serum phospholipids and carotid intima-media thickness in Spanish subjects with primary dyslipidemia. *Am J Clin Nutr*, United States 2010;92:186–93.
- [14] Yang K, Yang W, Mariadason J, Velcich A, Lipkin M, Augenlicht L. Dietary components modify gene expression: implications for carcinogenesis. *J Nutr* 2005;135:2710–4.
- [15] Hamer M, Steptoe A. Influence of specific nutrients on progression of atherosclerosis, vascular function, haemostasis and inflammation in coronary heart disease patients: a systematic review. *Br J Nutr* 2006;95:849–59.
- [16] Kark JD, Kaufmann NA, Binka F, Goldberger N, Berry EM. Adipose tissue n-6 fatty acids and acute myocardial infarction in a population consuming a diet high in polyunsaturated fatty acids. *Am J Clin Nutr* 2003;77:796–802.
- [17] Williams ES, Baylin A, Campos H. Adipose tissue arachidonic acid and the metabolic syndrome in Costa Rican adults. *Clin Nutr* 2007;26:474–82.
- [18] Koletzko B, Lien E, Agostoni C, Bohles H, Campoy C, Cetin I, et al. The roles of long-chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendations. *J Perinat Med* 2008;36:5–14.
- [19] Allayee H, Roth N, Hodis HN. Polyunsaturated fatty acids and cardiovascular disease: implications for nutrigenetics. *J Nutrigenet Nutrigenom* 2009;2:140–8.
- [20] Carlson SE. Docosahexaenoic acid supplementation in pregnancy and lactation. *Am J Clin Nutr* 2009;89:678S–84S.
- [21] Decsi T, Molnar D, Koletzko B. The effect of under- and overnutrition on essential fatty acid metabolism in childhood. *Eur J Clin Nutr* 1998;52:541–8.
- [22] de Groot RH, van Boxtel MP, Schiepers OJ, Hornstra G, Jolles J. Age dependence of plasma phospholipid fatty acid levels: potential role of linoleic acid in the age-associated increase in docosahexaenoic acid and eicosapentaenoic acid concentrations. *Br J Nutr* 2009;102:1058–64.

- [23] Fekete K, Marosvolgyi T, Jakobik V, Decsi T. Methods of assessment of n-3 long-chain polyunsaturated fatty acid status in humans: a systematic review. *Am J Clin Nutr* 2009;89:2070S–84S.
- [24] Al MD, van Houwelingen AC, Hornstra G. long-chain polyunsaturated fatty acids pregnancy pregnancy outcome. *Am J Clin Nutr* 2000;71:285S–91S.
- [25] Vlaardingerbroek H, Hornstra G. Essential fatty acids in erythrocyte phospholipids during pregnancy and at delivery in mothers and their neonates: comparison with plasma phospholipids. *Prostaglandins Leukot Essent Fatty Acids* 2004;71:363–74.
- [26] de Groot RH, Hornstra G, van Houwelingen AC, Roumen F. Effect of alpha-linolenic acid supplementation during pregnancy on maternal and neonatal polyunsaturated fatty acid status and pregnancy outcome. *Am J Clin Nutr* 2004;79:251–60.
- [27] Thorsdottir I, Gunnarsdottir I, Palsson GI. Birth weight, growth and feeding in infancy: relation to serum lipid concentration in 12-month-old infants. *Eur J Clin Nutr* 2003;57:1479–85.
- [28] Barker DJ, Martyn CN, Osmond C, Hales CN, Fall CH. Growth in utero and serum cholesterol concentrations in adult life. *BMJ* 1993;307:1524–7.
- [29] Iliescu C, Beghin L, Maes L, De Bourdeaudhuij I, Libersa C, Vereecken C, et al. Socioeconomic questionnaire and clinical assessment in the HELENA Cross-Sectional Study: methodology. *Int J Obes (London)* 2008;32Suppl5:S19–25.
- [30] Nagy E, Vicente-Rodriguez G, Manios Y, Beghin L, Iliescu C, Censi L, et al. Harmonization process and reliability assessment of anthropometric measurements in a multicenter study in adolescents. *Int J Obes (London)* 2008;32Suppl5:S58–65.
- [31] Tanner JM, Whitehouse RH. Clinical longitudinal standards for height, weight, height velocity, weight velocity, and stages of puberty. *Archives Dis Childhood* 1976;51:170–9.
- [32] Gonzalez-Gross M, Breidenassel C, Gomez-Martinez S, Ferrari M, Beghin L, Spinneker A, et al. Sampling and processing of fresh blood samples within a European multicenter nutritional study: evaluation of biomarker stability during transport and storage. *Int J Obes (London)* 2008;32(Suppl. (5)):S66–75.
- [33] Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 1959;37:911–7.
- [34] Christophe A, Matthijs F. New method for the determination of the fatty acid pattern of serum lipid classes. *Clin Chim Acta* 1967;16:39–43.
- [35] Vereecken CA, De Henauw S, Maes L. Adolescents' food habits: results of the Health Behaviour in School-aged Children survey. *Br J Nutr* 2005;94:423–31.
- [36] Brussaard JH, Löwik MR, Steingrimsdóttir L, Møller A, Kearney J, De Henauw S, Becker W, EFCOSUM Group. A European food consumption survey method—conclusions and recommendations. *Eur J Clin Nutr* 2002;56:S89–94.
- [37] Rogers I, Emmett P, Ness A, Golding J. Maternal fish intake in late pregnancy and the frequency of low birth weight and intrauterine growth retardation in a cohort of British infants. *J Epidemiol Commun Health* 2004;58:486–92.
- [38] Dirix CE, Kester AD, Hornstra G. Associations between term birth dimensions and prenatal exposure to essential and trans fatty acids. *Early Hum Dev* 2009.
- [39] Emken EA, Adlof RO, Gulley RM. Dietary linoleic acid influences desaturation and acylation of deuterium-labeled linoleic and linolenic acids in young adult males. *Biochim Biophys Acta* 1994;1213:277–88.
- [40] Decsi T, Molnar D, Koletzko B. Long-chain polyunsaturated fatty acids in plasma lipids of obese children. *Lipids* 1996;31:305–11.
- [41] Scaglioni S, Verduci E, Salvioni M, Bruzzese MG, Radaelli G, Zetterstrom R, et al. Plasma long-chain fatty acids and the degree of obesity in Italian children. *Acta Paediatr* 2006;95:964–9.